



Short communication

Simultaneous detection of West Nile and Japanese encephalitis virus RNA by duplex TaqMan RT-PCR



Silvia C. Barros^a, Fernanda Ramos^a, Líbia Zé-Zé^b, Maria J. Alves^b, Teresa Fagulha^a, Margarida Duarte^a, Margarida Henriques^a, Tiago Luís^a, Miguel Fevereiro^{a,*}

^a INIAV, Rua General Morais Sarmento, 1500-311 Lisboa, Portugal

^b Center for Vectors and Infectious Diseases Research, National Institute of Health (CEVDI-INSA), Av. da Liberdade 5, 2965-575 Águas de Moura, Portugal

A B S T R A C T

Article history:

Received 23 January 2013

Received in revised form 5 July 2013

Accepted 12 July 2013

Available online 25 July 2013

Keywords:

WNV

JEV

Diagnosis

Duplex RT-PCR

West Nile virus (WNV) and Japanese encephalitis virus (JEV) are important mosquito-borne viruses of the *Flaviviridae* family, associated with encephalitis, mainly in humans and horses. WNV is also pathogen for many bird species. The incidence of human and animal WNV infections in Europe has risen, mostly in recent years, and JEV was detected in 2011 in mosquitoes collected in Italy and may emerge in Europe in the same way as other flaviviruses had emerged recently (USUTU and Bagaza virus) and should be regarded as a potential threat to public health.

Prompt identification and discrimination between WNV and JEV provides critical epidemiological data for prevalence studies and public and animal health management policies. Here we describe a quantitative one-step duplex TaqMan RT-PCR, targeting non-structural protein 2A gene (NS2A-qRT-PCR), based on only one primer pair and two probes for differential diagnosis of WNV and JEV. Also this assay enables the detection of both WNV lineages (WNV-1 and WNV-2).

To access the specificity of NS2A-qRT-PCR a panel of different arboviruses were used. The assay was shown to be specific for both WNV lineages (WNV-1 and WNV-2), WNV related Kunjin virus and JEV, since no cross-reactions were observed with other tested arboviruses. Sensitivity of the assay was determined using serial dilutions of in vitro-transcribed RNA from WNV and JEV. The duplex NS2A-qRT-PCR assay was shown to be very sensitive, being able to detect 10 copies of WNV and JEV RNA.

This assay is a suitable tool for the diagnosis of WNV and JEV, and provides a valuable addition to the methods currently available for routine diagnosis of these zoonoses and for surveillance studies.

© 2013 Elsevier B.V. All rights reserved.

West-Nile virus (WNV) and Japanese encephalitis virus (JEV) are zoonotic mosquito-transmitted flaviviruses belonging to the family *Flaviviridae*, which cause public health problems worldwide. They have a single-stranded positive-sense RNA genome which encodes 10 proteins, three structural (C, prM and E) and seven non-structural proteins (NS1, NS2A, NS2B, NS3, NS4A, NS4B, NS5).

Japanese encephalitis virus (JEV) causes acute encephalitis with a mortality rate up to one-third of infected patients, and nearly half of the survivors suffer neurological or mental sequelae (Vaughn and Hoke, 1992).

WNV is the most distributed arbovirus in the world (Gould et al., 2003; Kramer et al., 2008), while JEV is prevalent in Southeast Asia, Indonesia, Australia, Papua New Guinea, and Pakistan (Ghosh and Basu, 2009). In Europe, two recent reports were published concerning the detection of JEV in birds and mosquitoes collected in Italy (Ravanini et al., 2012; Platonov et al., 2012), raising the risk that JEV

may emerge in Europe, in the same way as other flaviviruses (Usutu and Bagaza virus) have emerged (Vásquez et al., 2011; Gamino et al., 2012).

Phylogenetic studies based in the envelope protein show that WNV can be separated into two major lineages. WNV lineage 1 (WNV-1) is the most widespread, and has been detected for decades in Europe. It was assumed that these viruses are particularly pathogenic for birds, humans and horses (Solomon et al., 2003). WNV lineage 2 (WNV-2) was thought to be confined to sub-Saharan Africa and Madagascar until it was first detected in Europe in 2004 (Bakonyi et al., 2005). Initially it was postulated that WNV-2 was non-pathogenic for horses (Lanciotti et al., 1999) however retrospective studies demonstrated that some variants of WNV-2 can also cause severe symptoms in horses and humans (Botha et al., 2008; Venter et al., 2009).

Real-time RT-PCR methods are essential tools in laboratory diagnosis. Several methods have been developed for the laboratory diagnosis of WNV infection (reviewed by De Filette et al., 2012). Many of these methods were initially designed to detect strains of WNV-1 (Buckley et al., 2003; Lanciotti et al., 2000) but not

* Corresponding author. Tel.: +351 217115288.

E-mail address: miguel.fevereiro@iniav.pt (M. Fevereiro).




Fig. 1. Standard curves of TaqMan uniplex and duplex NS2A-qRT-PCR assay generated from the Ct values obtained against the log of known WNV and JEV RNA copies (dilution series of 10^1 – 10^9 WNV copies RNA per reaction and 10^1 – 10^6 JEV copies RNA per reaction). The regression equations are indicated.

strains of lineage 2, found in Africa and more recently in Europe. However, novel real-time RT-PCRs have been published for detection and discrimination of WNV-1 and WNV-2 (Del Amo et al., 2013; Eiden et al., 2010; Jimenez-Clavero et al., 2006; Linke et al., 2007). Moreover, molecular methods to discriminate between strains of WNV and JEV have been reported previously, using RT-PCR complemented with restriction fragment length polymorphism (RFLP) analysis (Shirato et al., 2003), conventional RT-PCR (Yeh et al., 2010) and real-time RT-PCR based on the use of a single probe and primer sets for both viruses (Shirato et al., 2005). Compared to the conventional RT-PCR, real-time RT-PCR is more sensitive, specific, rapid, and allowing quantification of viral genome; thus progressively replacing the conventional RT-PCR (Mackay et al., 2002).

The aim of the present study was to develop a duplex one-step TaqMan quantitative real-time RT-PCR (qRT-PCR) assay targeting the NS2A gene (NS2A-qRT-PCR), based in only one primer pair and two probes, for rapid and specific detection of both WNV lineages and JEV. This assay was rendered quantitative by using known copy numbers of WNV and JEV RNA transcripts.

Specific primer pair and probes were designed manually based on alignment of WNV and JEV sequences retrieved from NCBI database (<http://www.ncbi.nlm.nih.gov/>) and potential mismatched bases were taken into consideration (Table 1).

The duplex NS2A-qRT-PCR assay was carried out in 25 μ l reactions using the one-step RT-PCR kit (Qiagen, Germany). Each reaction mixture contained final concentrations of 1 μ M of each primer, 0.2 μ M of each TaqMan probe, 1X one-step RT-PCR buffer, 0.4 mM dNTP mix, 1 μ l enzyme mix (including RT and hot-start Taq polymerase) and 1 μ l of RNA template. The reaction was performed using the following thermal cycling program: one cycle of RT at 50 °C for 30 min, one cycle at 95 °C for 15 min to activate the hot-start Taq polymerase, 50 cycles of amplification of cDNA with melting at 95 °C for 20 s, annealing at 50 °C for 30 s and elongation at 72 °C for 30 s. The negative control contained PCR-grade water instead of template RNA.

The specificity of NS2-qRT-PCR assay was evaluated by testing WNV and JEV specific probes against RNAs extracted from several arboviruses (Table 2). With the exception of Kunjin and JEV, provided as RNAs by the European reference laboratory (ANSES—Maisons-Alfort, France), all other viruses were grown in VeroE6 cells in minimum essential medium (MEM) (Invitrogen, USA) containing 10% fetal bovine serum (FBS) and an antibiotic-antimycotic mixture (Invitrogen, USA) at 37 °C in a humidified 5% CO₂ environment. Viral RNA was extracted using MagCore[®] viral nucleic acid extraction kit (RBCbioscience, Taiwan) according to the manufacturer's recommendations.

As a control to monitor the integrity and specificity of RNA, an in-house pan-FLAVI conventional RT-PCR targeting the NS3 region was also performed (primers: 5461F-ATGGATGARGCTCAYTTCAC and 5669R-GTKATCCATYCRATCCA) and the amplicons were subjected to sequence analysis to confirm their specificity.

The results demonstrated that the assay was completely specific for JEV and WNV RNA, detecting both lineages of WNV (WNV-1 and WNV-2) as well as the WNV-1b Kunjin virus (Table 2). No cross-reaction was observed with the other ten closely related flaviviruses nor with the CHIKV and Toscana arbovirus. Pan-FLAVI conventional RT-PCR assay yielded positive results for all flaviviruses tested, which was confirmed by sequence analysis. No amplification was detected when non-template control was used.

For sensitivity determination, in vitro-transcribed NS2A genes from WNV and JEV were used. A 136 bp portion of WNV-Eg 101 and JEV-Nakayama NS2A genes were amplified by NS2A primers. The amplicons were purified with the Qiaex II gel extraction kit (Qiagen, Germany) and cloned into the pCR2.1 vector using One Shot TOP10 chemically competent *Escherichia coli* (Invitrogen, USA). Plasmid was extracted using the plasmid midi purification kit (Qiagen, Germany). The BamHI restriction enzyme (Biolabs, UK) was used to linearize NS2A-pCR2.1 plasmids prior to in vitro transcription with the MAXIscript kit (Ambion, UK), according to the manufacturer's instructions. Transcribed RNA was treated with RNase-free DNase and purified with the DyeEx 2.0 Spin kit (Qiagen, Germany).

Table 1
Primer pair and TaqMan probes for WNV and JEV detection.

Primer/probe	Sequence (5' → 3')	Tm (°C)	Position ^{d,e}
NS2A-F	CCITTTTCAGY ^a TGGGCCTTCTG	59.5	3548–3568 ^d
NS2A-3R	CAGTGTA ^b GTV ^b ATR ^c CCCCCAA	60.8	3555–3575 ^e 3682–3662 ^d 3690–3670 ^e
WNVpNS2A-3612	FAM-AGCCAAGATCAGCATGCCAGC-TAMRA	61.8	3612–3632 ^d
JEVpNS2A-3628	RED-TGACCAATTCTCGGGTTTGGGG-BHQ	62.1	3628–3650 ^e

^{a,b,c} Degenerate nucleotides: Y = C or T; V = A, C or G; and R = A or G. Fluorophores: FAM (6-carboxyfluorescein) and RED (Texas red). Quenchers: TAMRA (6-carboxy-tetramethyl-rhodamine) and BHQ-1 (black hole dark quencher 1). ^{d,e} Numbering according to the sequences of WNV Eg101 (AF260968) and JEV (U15763) respectively.

Table 2
Summary of the viral strains used in this study and of the results of NS2A-qRT-PCR (WNV and JEV probes).

Family	Virus	GenBank accession no.	NS2A-qRT-PCR	
			WNV	JEV
Flaviviridae	WNV-Eg101-lineage1 (WNV-1a)	AF260968	+	–
Flaviviridae	WNV-Uganda B956-lineage2 (WNV-2)	AF394221	+	–
Flaviviridae	Kunjin virus (WNV-1b)	AY274505	+	–
Flaviviridae	JEV-Nakayama	EF571853	–	+
Flaviviridae	DENV-1-US/Hawaii/1944	EU848545	–	–
Flaviviridae	DENV-2-New Guinea C	M29095	–	–
Flaviviridae	DENV-3-H87	M93130	–	–
Flaviviridae	DENV-4-H241	AY947539	–	–
Flaviviridae	Zika virus- MR766	AY632535	–	–
Flaviviridae	YFV-strain Asabi	AY640589	–	–
Flaviviridae	Ntaya	AF13392	–	–
Flaviviridae	TBEV-strain Far East Sofjin	AF013399	–	–
Flaviviridae	USUTU-strain Meise H	JQ219843	–	–
Bunyaviridae	Toscana virus-ISS.Phl.3	NC.006319	–	–
Togaviridae	CHIKV-LR2006.OPY1	DQ443544	–	–

RNA transcripts were stored at -80°C until use. Additionally, a High Fidelity PCR Mix (Qiagen, Germany) was used with NS2A primers, according to the manufacturer's instructions, for testing the absence of DNA. RNA concentration was determined by spectrophotometry using a NanoDrop 1000 (Thermo Scientific, USA) by determining the average concentration from three measurements. RNA copy number was determined by the following formula:

$$\text{Amount (RNA copies}/\mu\text{l)} = \frac{\text{X(g}/\mu\text{l)RNA} \times 6.022 \times 10^{23}}{\text{nt transcript length} \times 340}$$

The in vitro derived T7 promoter transcript obtained was 249 nucleotide long, of which 136 bases correspond to virus-specific sequence. The detection threshold of NS2A-qRT-PCR was determined by using 10-fold serial dilutions of RNA standards in order to amplify from 10^6 to 1 (JEV) and 10^9 to 1 (WNV) copy per reaction. Reactions were repeated in duplicate for each sample and a sample was considered positive when viral RNA was detected in both replicas. Standard curves for each probe were generated by plotting Ct values per two replicates per standard dilution versus the logarithm of the RNA copy number to determine analytical sensitivity and efficiency of the assay. The Ct values should remain within the range of linearity of a standard curve with a minimum correlation coefficient of 0.98. The amplification efficiency (E) was calculated with the equation $E = [10^{(-1/k)} - 1]$, where (k) is the slope of the linear regression line (Pfaffl, 2001). An efficiency of 1 corresponds to 100% amplification efficiency.

The sensitivity of duplex qRT-PCR was compared with that of uniplex RT-PCR. WNV and JEV, duplex and uniplex RT-PCR were performed simultaneously with the same concentration of RNA template per dilution (Fig. 1). Based upon the real-time PCR standard curve, the correlation coefficient (R^2) and RT-PCR efficiency (E) were obtained. The WNV-probe showed similar limits of detection of 10 copies in uniplex and duplex reactions, with a R^2 value of 0.9994 and 0.9992, and an $E = 97\%$ and $E = 100\%$, respectively for uniplex and duplex NS2A-qRT-PCR assays. For JEV, 10 copies RNA could be detected in uniplex and duplex reactions with an R^2 of 0.9916 and 0.9918, and an $E = 94\%$ and $E = 100\%$, respectively. No obvious interference or inhibition was observed in duplex NS2-qRT-PCR when compared with uniplex assay.

In this study, a TaqMan duplex one-step RT-PCR was developed for specific detection of both WNV lineages and JEV in a single run. WNV and JEV RNA could be distinguished by the differently labeled probes and the fluorescence signals for WNV (FAM) and JEV (Texas Red) showed no crosstalk. This assay has the potential for use for clinical diagnosis and epidemiological surveillance. Since WNV-2 was introduced into Europe the detection of both WNV lineages

by RT-PCR is mandatory for a diagnostic laboratory. Past international proficiency studies on WNV indicated that some laboratories were unable to detect WNV-2 virus genome by their RT-PCR diagnostic assays (Linke et al., 2011). On the other hand, although no clinical cases of JEV have occurred, the presence of JEV sequences in mosquitoes collected in Italy suggests the virus may emerge in Europe as it has emerged in other continents, therefore laboratories must be prepared for its diagnosis.

The NS2A-qRT-PCR assay described above proved specific for WNV and JEV, as no cross-reaction against a panel of other 12 arboviruses was detected. This assay exhibited high sensitivity, being able to detect as few as 10 copies of WNV and JEV RNA. We already used this duplex NS2A-qRT-PCR with success in proficiency tests conducted by the European reference laboratory. In the present study, the high number of sequences used to select the primers provides an upgraded method for the detection of those viruses and reduces the probability of false-negative detection due to new viral mutations.

The duplex NS2-qRT-PCR assay developed in this study provides a sensitive and rapid tool that is a valuable addition to the methods currently available for routine diagnosis of WNV and JEV in horses, humans, birds or mosquitoes. This assay improves the current diagnostic capability and can be easily implemented for screening a large number of samples in a rapid, sensitive, and reproducible way, for clinical and epidemiological purposes.

References

- Bakonyi, T., Hubalek, Z., Rudolf, I., Nowotny, N., 2005. Novel flavivirus or new lineage of West Nile virus, central Europe. *Emerging Infectious Diseases* 11, 225–231.
- Botha, E.M., Markotter, W., Wolfaardt, M., Paweska, J.T., Swanepoel, R., Palacios, G., Nel, L.H., Venter, M., 2008. Genetic determinants of virulence in pathogenic lineage 2 West Nile virus strains. *Emerging Infectious Diseases* 14, 222–230.
- Buckley, A., Dawson, A., Moss, S.R., Hinsley, S.A., Bellamy, P.E., Gould, E.A., 2003. Serological evidence of West Nile virus, USUTU virus and Sindbis virus infection in birds in the UK. *Journal of General Virology* 84, 2807–2817.
- De Filette, M., Ulbert, S., Diamond, M., Sanders, N., 2012. Recent progress in West Nile virus diagnosis and vaccination. *Veterinary Research* 43, 16.
- Del Amo, J., Sotelo, E., Fernández-Pinero, J., Gallardo, C., Llorente, F., Agüero, M., Jiménez-Clavero, M.A., 2013. A novel quantitative multiplex real-time RT-PCR for the simultaneous detection and differentiation of West Nile virus lineages 1 and 2, and of USUTU virus. *Journal of Virological Methods* 189, 321–327.
- Eiden, M., Vina-Rodríguez, A., Hoffmann, B., Ziegler, U., Groschup, H.M., 2010. Two new real-time quantitative reverse transcription polymerase chain reaction assays with unique target sites for the specific and sensitive detection of lineages 1 and 2 West Nile virus strains. *Journal of Veterinary Diagnostic Investigation* 22, 748–753.
- Gamino, V., Gutiérrez-Guzmán, A.-V., Fernández-de-Mera, I.G., Ortíz, J.-A., Durán-Martín, M., de la Fuente, J., Gortázar, C., Höfle, U., 2012. Natural Bagaza virus infection in game birds in southern Spain. *Veterinary Research* 43, 65.
- Ghosh, D., Basu, A., 2009. Japanese encephalitis – a pathological and clinical perspective. *PLoS Neglected Tropical Diseases* 3, e437.

- Gould, E.A., de Lamballerie, X., Zanotto, P.M., Holmes, E.C., 2003. *Origins, evolution, and vector/host coadaptations within the genus flavivirus*. *Advances in Virus Research* 59, 277–314.
- Jimenez-Clavero, M.A., Aguero, M., Rojo, G., Gomez-Tejedor, C., 2006. *A new fluorogenic real-time RT-PCR assay for detection of lineage 1 and lineage 2 West Nile viruses*. *Journal of Veterinary Diagnostic Investigation* 18, 459–462.
- Kramer, L.D., Styer, L.M., Ebel, G.D., 2008. *A global perspective on the epidemiology of West Nile virus*. *Annual Review of Entomology* 53, 61–81.
- Lanciotti, R.S., Roehrig, J., Deubel, V., Smith, J., Parker, M., Steele, K., Crise, B., Volpe, K.E., Crabtree, M.B., Scherret, J.H., Hall, R.A., MacKenzie, J.S., Cropp, C.B., Panigrahy, B., Ostlund, E., Schmitt, B., Malkinson, M., Banet, C., Weissman, J., Komar, N., Savage, H.M., Stone, W., McNamara, T., Gubler, D.J., 1999. *Origin of the West Nile virus responsible for an outbreak of encephalitis in the northeastern United States*. *Science* 286, 2333–2337.
- Lanciotti, R.S., Kerst, A., Nasci, R., Godsey, M., Mitchell, C., Savage, H., Komar, N., Panella, N.A., Allen, B.C., Volpe, K.E., David, B.S., Roehrig, J.T., 2000. *Rapid detection of West Nile virus from human clinical specimens, field-collected mosquitoes, and avian samples by a TaqMan reverse transcriptase-PCR assay*. *Journal of Clinical Microbiology* 38, 4066–4071.
- Linke, S., Ellerbrok, H., Niedrig, M., Nitsche, A., Pauli, G., 2007. *Detection of West Nile virus lineages 1 and 2 by real-time PCR*. *Journal of Virological Methods* 146, 355–358.
- Linke, S., Mackay, W., Scott, C., Wallace, P., Niedrig, M., 2011. *Second external quality assessment of the molecular diagnostic of West Nile virus: are there improvements towards the detection of WNV?* *Journal of Clinical Virology* 52, 257–260.
- Mackay, I.M., Arden, K.E., Nitsche, A., 2002. *Real-time PCR in virology*. *Nucleic Acids Research* 30, 1292–1305.
- Pfaffl, M.W., 2001. *A new mathematical model for relative quantification in real-time RT-PCR*. *Nucleic Acids Research* 29, e45.
- Platonov, A., Rossi, G., Karan, L., Mironov, K., Busani, L., Rezza, G., 2012. *Does the Japanese encephalitis virus (JEV) represent a threat for human health in Europe? Detection of JEV RNA sequences in birds collected in Italy*. *Euro Surveillance* 17 (32), pii:20241.
- Ravanini, P., Huhtamo, E., Ilaria, V., Crobu, M.G., Nicosia, A.M., Servino, L., Rivasi, F., Allegrini, S., Miglio, U., Magri, A., Minisini, R., Vapalahti, O., Boldorini, R., 2012. *Japanese encephalitis virus RNA detected in Culex pipiens mosquitoes in Italy*. *Eurosurveillance* 17, 2.
- Shirato, K., Mizutani, T., Kariwa, H., Takashima, I., 2003. *Discrimination of West Nile virus and Japanese encephalitis virus strains using RT-PCR RFLP analysis*. *Microbiology and Immunology* 47, 439–445.
- Shirato, K., Miyoshi, H., Kariwa, H., Takashima, I., 2005. *Detection of West Nile virus and Japanese encephalitis virus using real-time PCR with a probe common to both viruses*. *Journal of Virological Methods* 126, 119–125.
- Solomon, T., Ooi, M.H., Beasley, D.W., Mallewa, M., 2003. *West Nile encephalitis*. *BMJ* 326, 865–869.
- Vásquez, A., Jiménez-Clavero, M.A., Franco, L., Donoso-Mantke, O., Sambri, V., Niedrig, M., Zeller, H., Tenorio, A., 2011. *Usutu virus-potential risk of human disease in Europe*. *Eurosurveillance* 16 (31), pii:19935.
- Vaughn, D.W., Hoke Jr., C.H., 1992. *The epidemiology of Japanese encephalitis: prospects for prevention*. *Epidemiologic Reviews* 14, 197–221.
- Venter, M., Human, S., Zaayman, D., Gerdes, G.H., Williams, J., Steyl, J., Leman, P.A., Paweska, J.T., Setzkorn, H., Rous, G., Murray, S., Parker, R., Donellan, C., Swanepoel, R., 2009. *Lineage 2 West Nile virus as cause of fatal neurologic disease in horses*. *South Africa – Emerging infectious diseases* 15, 877–884.
- Yeh, J.Y., Lee, J.H., Seo, H.J., Park, J.Y., Moon, J.S., Cho, I.S., Lee, J.B., Park, S.Y., Song, C.S., Choi, I.S., 2010. *Fast duplex one-step reverse transcriptase PCR for rapid differential detection of West Nile and Japanese encephalitis viruses*. *Journal of Clinical Microbiology* 48, 4010–4014.